

## ARTICLE INFORMATION

Received: February, 27, 2024

Revised: April, 03, 2024

Available online: April, 17, 2024

at : <https://ejournal.malahayati.ac.id/index.php/minh>

## Maternal knowledge regarding antenatal care services, history of exclusive breastfeeding and stunting among toddlers in West Java, Indonesia

Astri Mufti Aini\*, Maria Komariah, Lilis Mamuroh, Laili Rahayuwati, Yanti Hermayanti, Ristina Mirwanti, Theresia Eriyani

Fakultas Keperawatan Universitas Padjadjaran

Corresponding author: \*E-mail: [astri20006@mail.unpad.ac.id](mailto:astri20006@mail.unpad.ac.id)

### Abstract

**Background:** A mother's knowledge of ANC can increase interest and motivation in carrying out pregnancy checks which aim to detect problems early. Exclusive breastfeeding is one effort that can be made to meet nutritional needs to prevent stunting.

**Purpose:** Analyzing the correlation between maternal knowledge of ANC, exclusive breastfeeding history, and stunting incidence in Hamlet III, Sukamulya Village.

**Method:** Quantitative correlational, with a population of 68 families of stunted toddlers living in Hamlet III, Sukamulya Village. The sample size used for the research is total sampling. Secondary data is used for the research with instruments that include a combination of "Environmental Modification through Sanitation, Clean Water, Hygiene and Nutrition for Stunting Prevention" with the research instruments of Khairani, (2012) and Fitriani, (2021). The data analysis method used is univariate and bivariate analysis (Spearman rho).

**Results:** There was no significant relationship between ANC knowledge and the incidence of stunting, obtained  $r=0.114$  and  $p=0.353$ . Likewise, exclusive breastfeeding also did not show a significant relationship with the incidence of stunting, obtained  $r=0.007$  and  $p=0.955$ .

**Conclusion:** There is no relationship between maternal ANC knowledge and exclusive breastfeeding with the incidence of stunting.

**Keywords:** Antenatal Check-Up; Exclusive Breastfeeding; Incidence of Stunting; Level of Knowledge; Mother.

### INTRODUCTION

Stunting is a condition where a child's growth and development fail, which is caused by malnutrition over a long period (Magasida & Erawati, 2022). The World Health Organization (WHO) defines stunting as a condition where the z score for height for age is  $< -2$  SD. Factors that cause stunting can come from various dimensions (multidimensional) and can be broken down into parent, child and household environmental factors (Sitanggang Kasim, & Sari, 2020).

The prevalence of stunting in Indonesia is relatively high. Indonesia is ranked 10th in Southeast

Asia. The prevalence of stunting in Indonesia in 2022 will be 21.6% (Ministry of Health of the Republic of Indonesia, 2023). If the incidence of stunting reaches 20% or more, it can be said to be a public health problem (Apriluana & Fikawati, 2018). Based on this, it can be concluded that Indonesia has still not succeeded in overcoming the stunting problem and requires efforts to overcome it.

Prevention of stunting can be started during pregnancy, one of which is by regularly checking the condition of the pregnancy. Antenatal care is important for the early detection of problems during

pregnancy and for maintaining the health of both the mother and the baby (Dharmayanti, Azhar, Tjandrarini, & Hidayangsih, 2019). According to Health Personnel Education Center, neglecting antenatal care can lead to insufficient knowledge about proper methods of pregnancy care, and failure to detect pregnancy problems promptly (Dewanggayastuti, Surinati, & Hartati, 2022). This includes danger signs of pregnancy, signs of childbirth complications, comorbidities, and other complications that can potentially trigger bleeding during childbirth and even lead to maternal death. One of the examinations carried out during antenatal care is a weight examination, the aim of which is to show possible growth disorders in the fetus (Wiyandani, 2019). One of the effects of poor fetal development during pregnancy is the low birth weight (LBW), which if left for a long time has the potential to cause stunting.

The execution of antenatal care is influenced by various factors. Health behavior is influenced by driving, supporting, and reinforcing factors, with knowledge being the initial factor in forming health behavior (Green, 2001; Asmarasari, & Astuti, 2019). Individuals with higher levels of knowledge tend to exhibit better health behavior (Hermawan, Yani, Yulianita, & Rahayuwati, 2023). That is accordance that, a mother's knowledge about antenatal care can enhance her interest and motivation in carrying out antenatal care (Panjaitan, Santosa, & Utama, 2019). One research shows that babies born to mothers who don't receive standardized ANC have 2.28 times higher risk of stunting than those born to mothers who do (Amini, 2016).

In addition to irregular pregnancy checks, malnutrition in the first 1000 days of life can also be a cause of stunting. Actions that can be taken after labor are optimization of early breastfeeding initiation practices, followed by exclusive breastfeeding for 6 months without being accompanied by giving fluids or other foods (Rahmadi, 2016). There is a significant relationship between exclusive breastfeeding and stunting incidence with a p-value of 0.000 (Sampe, Toban, & Madi, 2020). This study was supported by research which found a significant relationship between exclusive breastfeeding and stunting ( $p = 0.001$ ) (Pratama, & Irwandi, 2021).

Contrary to previous research, obtained a  $p$  value = 1.001 which shows that there is no relationship between exclusive breastfeeding and the incidence of stunting (Malka, Musni, & Fatimah, 2020). The things that affect the incidence of stunting in this study are early pregnancy and ANC visits. In the other study, there was no correlation between stunting and exclusive breastfeeding (Hikmahrachim, Rohsiswatmo, & Ronoatmodjo, 2020). The study identified maternal age, socioeconomics, education level, and quality of ANC as factors contributing to stunting.

Sukamulya Village is one of the stunting locus villages in Bandung Regency with a total stunting incidence of 68 toddlers (33.6%) out of a total of 202 toddlers residing in Hamlet III of Sukamulya Village. This still shows that the target of the West Java Provincial Government to become a zerostunting province in 2023 has not been achieved. In addition, the average education level of the local productive age population is graduating from elementary school to junior high school. Education level can impact knowledge and information acceptance, so higher education levels lead to better understanding (Angraini, Amin, Pratiwi, Febriawati, & Yanuarti, 2021). This may be one of the factors causing stunting in Hamlet III of Sukamulya Village.

Seeing the low level of education of the local community and the high incidence of stunting, encouraged researchers to further examine how maternal knowledge about ANC and exclusive breastfeeding can affect stunting incidence in Hamlet III of Sukamulya Village. The lack of literature regarding the correlation between ANC knowledge and stunting incidence prompted further research. In addition, differences in previous studies on the correlation between exclusive breastfeeding and stunting incidence also encourages researchers to examine the problem more deeply.

## RESEARCH METHOD

This study is a form of secondary research that uses correlational quantitative research design. The population used is the population in the primary study, namely families who have stunting toddlers and live in Hamlet III of Sukamulya Village, Rancaekek District, Bandung Regency. The sampling technique used is total sampling, with inclusion criteria for mothers who

**Astri Mufti Aini\***, Maria Komariah, Lilis Mamuroh, Laili Rahayuwati, Yanti Hermayanti, Ristina Mirwanti, Theresia Eriyani

Fakultas Keperawatan Universitas Padjadjaran  
Corresponding author: \*E-mail: astri20006@mail.unpad.ac.id

DOI: <https://doi.org/10.33024/minh.v7i2.226>

Maternal knowledge regarding antenatal care services, history of exclusive breastfeeding and stunting among toddlers in West Java, Indonesia

have at least 1 or more stunting toddlers, mothers who have given birth within the last 5 years, have a history of ANC, and provide complete data needed by researchers. The primary study included 68 mothers with stunted toddlers.

The instrument used is a combination of primary research instruments "Environmental Modification Through Sanitation, Clean Water, Hygiene, and Nutrition for Stunting Prevention" developed by Rahayuwati, et al. (2023) with research instruments used by (Khairani, 2012) and (Fitriani, 2021). The results of the validity test show that the result of r count is greater than r table, so it is said that the instrument used is valid. The results of the reliability test show an alpha value greater than 0.7 so that the questionnaire can be said to be reliable and can be used. ANC Knowledge Level is categorized as good with a value of 76-100%; Sufficient if 56-75%; and

Less if <55%. incidence of stunting are categorized as moderate with height for age with a value -3 SD to -2 SD; and severe if height for age < -3 SD.

This research uses secondary data and has received from the Research Ethics Committee of Padjadjaran University with ethical number 37/UN6. KEP/EC/2023.

After obtaining the secondary data, it was analyzed using both univariate and bivariate analysis. The data obtained was analyzed using univariate analysis to observe the frequency distribution. Bivariate analysis was conducted to examine the correlation between ANC knowledge, exclusive breastfeeding, and incidence of stunting. To test the correlation between the variables studied with the available data, the statistical test used is Spearman's rho.

**RESEARCH RESULTS**

**Table 1. Characteristics of Respondents (N = 68)**

Variables	Results
<b>Maternal Age (Mean±SD)(Range)(Year)</b>	(28.17±8.725)(19-49)
<b>Education Level (n/%)</b>	
Elementary School	19/27.9
Junior High School	37/54.4
Senior High School	12/17.7
<b>Number of Children (n/%)</b>	
1 child	24/35.3
2 children	31/45.6
3 children	7/10.3
≥ 4 children	6/8.8
<b>Revenue per Month (n/%)</b>	
Under Wages City/Regency	50/73.5
Wages equivalent – Rp. 5.000.000	18/26.5
<b>ANC Knowledge Level (n/%)</b>	
Good	21/30.9
Sufficient	38/55.9
Less	9/13.2
<b>Exclusive Breastfeeding (n/%)</b>	
≥ 6 Month	32/47.1
< 6 Month	36/52.9
<b>Incidence of Stunting (n/%)</b>	
Moderate	38/55.9
Severe	30/44.1

Astri Mufti Aini\*, Maria Komariah, Lilis Mamuroh, Laili Rahayuwati, Yanti Hermayanti, Ristina Mirwanti, Theresia Eriyani

Fakultas Keperawatan Universitas Padjadjaran  
Corresponding author: \*E-mail: astri20006@mail.unpad.ac.id

DOI: <https://doi.org/10.33024/minh.v7i2.226>

## Maternal knowledge regarding antenatal care services, history of exclusive breastfeeding and stunting among toddlers in West Java, Indonesia

Based on the data presented in Table 1, the mean maternal age is 28.17 with a standard deviation of 8.725 and an age range between 19 and 49 years. In terms of education, the majority of mothers had completed junior high school education, namely 37 (54.4%). In addition, most mothers had two children, namely 31 (45.6%). Most families have a monthly income below the Regency/City wage, which is 50 (73.5%). The majority of mothers' level of knowledge about antenatal care (ANC) was in the sufficient category, namely 38 (55.9%). The majority of toddlers do not receive exclusive breast milk (< 6 months), namely 36 (52.9%), and the majority of stunting incidents are in the moderate stunting category, namely 38 (55.9%).

**Table 2. Correlation Analysis (N = 68)**

Variable	Incidence of Stunting		$\rho$	r
	Moderate (n=38)	Severe (n=30)		
<b>ANC Knowledge Level (n/%)</b>				
Good	14/36.8	7/23.3	0.353	0.114
Sufficient	19/50.0	19/63.3		
Less	5/13.2	4/13.4		
<b>Exclusive Breastfeeding (n/%)</b>				
≥ 6 Month	18/47.4	14/46.7	0.955	0.007
< 6 Month	20/52.6	16/53.3		

Table 2 shows that the majority of mothers who have sufficient knowledge have toddlers in the moderate stunting category, namely 19 cases (50.0%). Statistical analysis using the Spearman rho test produces  $r = 0.114$  and  $p = 0.353$ . and the majority of babies who do not receive exclusive breastfeeding suffer from moderate stunting, namely around 20 (52.6%). The results of statistical tests with Spearman's Rho show a correlation coefficient  $r = 0.007$  and p-value  $p = 0.955$ .

**DISCUSSION****ANC Knowledge and Stunting**

This study involved 68 respondents who were mothers who had *stunted* children aged 5 years and under and lived in Hamlet III of Sukamulya Village. Out of 68 respondents, it was discovered that 21 mothers (30.9%) had a good understanding of ANC. Out of 68 mothers, 38 (55.9%) had sufficient knowledge about ANC. Only 13.2% of mothers have a level of knowledge of ANC that is below 9.

The data obtained was then processed and carried out statistical analysis using *Spearman rho* where the results of the correlation coefficient of 0.114 and sig value were obtained. amounted to 0.353. Based on the statistical analysis, there is no significant correlation between maternal knowledge about antenatal care (ANC) and the incidence of stunting. This may happen because knowledge about ANC is not a direct factor causing stunting, but an indirect factor.

Based on research at Ridos Medan Hospital, of the 89 samples, 56 (62.9%) had sufficient knowledge

and 23 (30.3%) had good knowledge. Research also shows that good knowledge can increase mothers' interest in attending antenatal care (ANC) visits (Panjaitan et al., 2019). This is also supported by research conducted at the Abiansemal I Community Health Center, where from a total of 32 samples, 17 (53.1%) respondents had good knowledge, 13 (40.6%) had sufficient knowledge, while 2 (6.3%) respondents have less knowledge. Based on the results of bivariate analysis, there is a relationship between knowledge and ANC regularity, which is indicated by a p-value of 0.000 and a coefficient of  $r = 0.876$  (Padesi et al., 2021).

However, knowledge is not a direct factor causing stunting, because in addition to knowledge, experience can also be one of the influencing factors. Mothers who have had more than one pregnancy have greater knowledge compared to first-time mothers (Kanu, Tang, & Liu, 2014). This is because mothers who have been pregnant multiple

**Astri Mufti Aini\***, Maria Komariah, Lilis Mamuroh, Laili Rahayuwati, Yanti Hermayanti, Ristina Mirwanti, Theresia Eriyani

Fakultas Keperawatan Universitas Padjadjaran  
Corresponding author: \*E-mail: astri20006@mail.unpad.ac.id

DOI: <https://doi.org/10.33024/minh.v7i2.226>

times have a history of more frequent pregnancy checks than first-time mothers.

Thus, it can be said that antenatal care (ANC) visits are more appropriate if they are said to be a factor causing stunting, while ANC knowledge is only an indirect factor. The research conducted by Amini, (2016) with a total sample of 128 respondents, showed that non-standardized ANC visits were 57.8% and more common in stunted toddlers compared to non-stunted toddlers. Chi-square analysis indicated a significant association between stunting in toddlers aged 12-59 months and their ANC visits. The study found that mothers who did not receive standardized antenatal care had a 2.28 times higher risk of having stunted toddlers compared to those who did.

### Breastfeeding and Stunting

This study aimed to examine the history of exclusive breastfeeding among 68 mothers living in Hamlet III of Sukamulya Village. According to the study, 32 infants (47.1%) at Hamlet III in Sukamulya Village were exclusively breastfed. Meanwhile, 36 (52.9%) toddlers do not get exclusive breastfeeding.

The data obtained were then processed and statistical analysis was carried out using *Spearman rho*. The results obtained are a correlation coefficient of 0.007 and a sig value. amounted to 0.955. From the results of the statistical analysis it can be concluded that there is no significant correlation between exclusive breastfeeding and the stunting incidence.

Breast milk is a liquid produced by the breast glands that contains essential nutrients for babies. Exclusive breastfeeding for 6 months without being followed by fluids or other foods such as formula milk, water, honey, bananas, porridge, and others. Exclusive breastfeeding for 6 months without being followed by other foods can help children get adequate nutritional intake and the baby's intestines are able to digest incoming intake optimally.

The results of this study contradict the results of research in Buntu Malangka District, Mamasa Regency, which found a correlation between exclusive breastfeeding and the incidence of stunting in toddlers. Where toddlers who do not receive exclusive breastfeeding for 6 months have a 61 times higher risk of experiencing stunting than toddlers who receive exclusive breastfeeding (Sampe, et al., 2020). This is also supported by the results of research at the

Hinai Kiri Health Center, Secanggang District, Langkat Regency, on 60 samples which showed a significant relationship between exclusive breastfeeding and the incidence of stunting. Where it is said that toddlers who receive exclusive breast milk have 0.5 times higher protection in preventing stunting (Pratama & Irwandi, 2021).

However, several studies show that exclusive breastfeeding is not significantly related to the incidence of stunting. Research conducted through a survey of 162 respondents showed that there was no relationship between exclusive breastfeeding and the incidence of stunting (Hikmahrachim et al., 2020). This is supported by research in Bone District, with p-value = 0.892 which shows that there is no relationship between exclusive breastfeeding and the incidence of stunting (Malka et al., (2020). The results of other studies show that the factors associated with the incidence of stunting are maternal age during pregnancy (p=0.003) and antenatal care (ANC) (p=0.033) (Mutunga, Frison, Rava, & Bahwere, 2020; Appiah, Amu, Osei, Konlan, Mumuni, Verner, & Kim, 2021; Lengkong, Wagey, & Tatura, 2023).

### Incidence of Stunting

Stunting is a condition where a child's growth and development fails which can be caused by poor nutrition, repeated infections and inadequate psychosocial stimulation. Stunting is a condition where the z score for height for age is < -2 SD. This event is a long-term impact that occurs from pregnancy to the first 1000 days of life and only appears at the age of 2 years.

This study involved 68 respondents who lived in Hamlet III of Sukamulya Village and had stunting toddlers. From the results of the study, data were obtained that there were 38 (55.9%) stunted toddlers with moderate stunting category. While 30 (44.1%) other toddlers are stunted in the category of severe stunting.

The incidence of stunting can be influenced by various factors (multidimensional), including poor parenting practices such as breastfeeding and complementary foods, limited access to health services so that pregnancy checks (ANC) and postnatal care cannot be fulfilled properly, access to nutritious food, clean water, and poor environmental sanitation, and chronic lack of energy

Astri Mufti Aini\*, Maria Komariah, Lilis Mamuroh, Laili Rahayuwati, Yanti Hermayanti, Ristina Mirwanti, Theresia Eriyani

Fakultas Keperawatan Universitas Padjadjaran  
Corresponding author: \*E-mail: astri20006@mail.unpad.ac.id

DOI: <https://doi.org/10.33024/minh.v7i2.226>

Maternal knowledge regarding antenatal care services, history of exclusive breastfeeding and stunting among toddlers in West Java, Indonesia

(SEZ) conditions in mothers. In addition to these things, other factors such as socioeconomic status, education level, premature birth, short birth length, and maternal age and height can also be factors that may affect the incidence of stunting. Therefore, further studies are needed on factors that affect the incidence of stunting, considering that stunting risk factors can occur from various factors.

The incidence of stunting greatly impacts the condition of babies in the future, such as susceptible to disease, impaired brain and physical development, non-optimal posture, increased risk of obesity and comorbidities, and decreased learning achievement. For this reason, an effort is needed to prevent stunting, starting from pregnancy to the future. Starting from maintaining health during pregnancy by consuming nutritious foods and routinely conducting pregnancy checks, conducting exclusive breastfeeding for 6 months and providing complementary foods after the age of 6 months, always monitoring the growth and development of toddlers at Integrated Healthcare Center, and increasing access to clean water and sanitation facilities. Thus, it is hoped that the incidence of stunting can be reduced, and the target of the West Java Provincial government in realizing West Java as a zero stunting province can be realized.

## CONCLUSION

Research of 68 respondents from families with infants with stunting toddlers in Hamlet III of Sukamulya Village, found that mothers' knowledge of antenatal care (ANC) and exclusive breastfeeding there was no correlation with the stunting incidence. Since there are many factors (multidimensional), that contribute to stunting, further research is needed to examine the factors that cause stunting in Hamlet III of Sukamulya Village.

## REFERENCES

- Amini, A. (2016). Hubungan Antenatal Care (ANC) dengan Kejadian Stunting pada Balita Usia 12-59 Bulan di Kabupaten Lombok Utara Provinsi NTB Tahun 2016. Universitas Aisyiyah Yogyakarta, 2–22. file:///C:/Users/Acer/Downloads/anc.pdf
- Angraini, W., Amin, M., Pratiwi, B. A., Febriawati, H., & Yanuarti, R. (2021). Pengetahuan Ibu, Akses Air

Bersih dan Diare dengan Stunting di Puskesmas Aturan Mumpo Bengkulu Tengah. *Jurnal Kesehatan Masyarakat Khatulistiwa*, 8(2), 92–102.

- Appiah, P. K., Amu, H., Osei, E., Konlan, K. D., Mumuni, I. H., Verner, O. N., & Kim, S. Y. (2021). Breastfeeding and weaning practices among mothers in Ghana: A population-based cross-sectional study. *Plos one*, 16(11), e0259442.
- Apriluana, G., & Fikawati, S. (2018). Analisis Faktor-Faktor Risiko terhadap Kejadian Stunting pada Balita (0-59 Bulan) di Negara Berkembang dan Asia Tenggara. *Media Litbangkes*, Vol. 28(4), 247–256. <https://doi.org/https://doi.org/10.22435/mpk.v28i4.472>
- Asmarasari, B., & Astuti, R. S., (2019). Analisis Perilaku Ibu dalam Memberikan ASI Eksklusif di Kota Semarang (Studi Kasus Puskesmas Genuk). *Journal of Public Policy and Management Review*, 8(4), 267–279.
- Dewanggayastuti, K. I., Surinati, I. D. A. K., & Hartati, N. N. (2022). Kepatuhan Ibu Hamil melakukan Kunjungan Antenatal Care (ANC) Pada Masa Pandemi Covid-19. *Jurnal Gema Keperawatan*, 15(1), 54–67.
- Dharmayanti, I., Azhar, K., Tjandrarini, D. H., & Hidayangsih, P. S. (2019). Pelayanan Pemeriksaan Kehamilan Berkualitas Yang Dimanfaatkan Ibu Hamil untuk persiapan Persalinan di Indonesia. *Jurnal Ekologi Kesehatan*, 18(1), 60–69. <https://doi.org/10.22435/jek.18.1.1777.60-69>
- Fitriani, L. (2021). Hubungan Pemberian ASI Eksklusif Dengan Kejadian Stunting Pada Balita Di Puskesmas Singkuang Tahun 2021. Skripsi, Universitas Aufa Royhan.
- Green, L. W. (2001). From research to “best practices” in other settings and populations. *American journal of health behavior*, 25(3), 165-178.
- Hermawan, S. I., Yani, D. I., Yulianita, H., & Rahayuwati, L. (2023). Correlation Mothers' Knowledge About Stunting and Exclusive Breastfeeding Behavior. *Journal of Nursing Care*, 6(2), 9.
- Hikmahrachim, H. G., Rohsiswatmo, R., & Ronoatmodjo, S. (2020). Efek ASI Eksklusif

**Astri Mufti Aini\***, Maria Komariah, Lilis Mamuroh, Laili Rahayuwati, Yanti Hermayanti, Ristina Mirwanti, Theresia Eriyani

Fakultas Keperawatan Universitas Padjadjaran  
Corresponding author: \*E-mail: astri20006@mail.unpad.ac.id

DOI: <https://doi.org/10.33024/minh.v7i2.226>

Maternal knowledge regarding antenatal care services, history of exclusive breastfeeding and stunting among toddlers in West Java, Indonesia

- terhadap Stunting pada Anak Usia 6-59 bulan di Kabupaten Bogor tahun 2019. *Jurnal Epidemiologi Kesehatan Indonesia*, 3(2), 77-82. <https://doi.org/10.7454/epidkes.v3i2.3425>
- Kanu, J. S., Tang, Y., & Liu, Y. (2014). Assessment on the knowledge and reported practices of women on maternal and child health in rural Sierra Leone: A cross-sectional survey. *PLoS ONE*, 9(8). <https://doi.org/10.1371/journalone.0105936>
- Khairani, L. (2012). Hubungan Pengetahuan Dan Sikap Ibu Hamil Terhadap Kunjungan Pemeriksaan Kehamilan Di Wilayah Kerja Puskesmas 4 Ulu Palembang.
- Lengkong, M., Wagey, F. W., & Tatura, S. N. (2023). Analysis Of Factors Affecting The Incidence Of High-Risk Pregnancies In The Working Area Of The Mubune Health Center, Northern Minahasa District. *Best Journal of Administration and Management*, 2(1), 14-26.
- Magasida, D., & Erawati, E. (2022). Hubungan Antenatal Care Terhadap Kejadian Stunting Pada Balita Usia 24-59 Bulan. *Midwifery Research*, 11(1).
- Malka, S., Musni, M., & Fatimah, S. (2020). Kehamilan dini, antenatal care, ASI eksklusif dan pengetahuan gizi terhadap stunting pada balita. *J Kebidanan*, 7(1), 59-64. <https://scholar.archive.org/work/6llu6ufw3zdblammv7y6qomgte/access/wayback/http://ejournalmalahayati.ac.id/index.php/kebidanan/article/download/3364/pdf>
- Ministry of Health of the Republic of Indonesia. (2023). Prevalensi Stunting di Indonesia Turun ke 21,6% dari 24,4%. Accessed from: <https://sehatnegeriku.kemkes.go.id/baca/rilis-media/20230125/3142280/prevalensi-stunting-di-indonesia-turun-ke-216-dari-244/>
- Mutunga, M., Frison, S., Rava, M., & Bahwere, P. (2020). The forgotten agenda of wasting in Southeast Asia: Burden, determinants and overlap with stunting: A review of nationally representative cross-sectional demographic and health surveys in six countries. *Nutrients*, 12(2), 559.
- Padesi, N. L. W., Suamiti, N. W., & Sriasih, N. G. K. (2021). Hubungan Pengetahuan Tentang Kunjungan Antenatal Care Dengan Keteraturan Kunjungan Antenatal Care Ibu Hamil Trimester Iii Di Masa Pandemi Covid-19. *Jurnal Ilmiah Kebidanan (The Journal Of Midwifery)*, 9(2), 183-189. <https://doi.org/10.33992/jik.v9i2.1421>
- Panjaitan, P. R. M., Santosa, H., & Utama, S. (2019). Associations between Education, Knowledge, Attitude, and Maternal Intention on Antenatal Care Visit. *Journal of Maternal and Child Health*, 4(6), 516-521. <https://doi.org/10.26911/thejmch.2019.04.06.12>
- Pratama, M. R., & Irwandi, S. (2021). Hubungan Pemberian Asi Eksklusif Dengan Stunting Di Puskesmas Hinai Kiri, Kecamatan Secanggang, Kabupaten Langkat. *Jurnal Kedokteran STM (Sains Dan Teknologi Medik)*, 4(1), 17-25. <https://doi.org/10.30743/stm.v4i1.65>
- Rahayuwati, L., Ibrahim, K., Hendrawati, S., Sari, C. W. M., Yani, D. I., Pertiwi, A. S. P., & Fauziyyah, R. N. P. (2022). Pencegahan Stunting melalui Air Bersih, Sanitasi, dan Nutrisi. *Warta LPM*, 356-365.
- Rahmadi, A. (2016). Hubungan Antara Pengetahuan Tentang Antropometri Dengan Ketidakpuasan Terhadap Body Image. *Jurnal Kesehatan Metro Sai Wawai*, VII(2), 50-56.
- Sampe, A., Toban, R. C., & Madi, M. A. (2020). Hubungan Pemberian ASI Eksklusif dengan Kejadian Stunting pada Balita. *Jurnal Ilmiah Kesehatan Sandi Husada*, 11(1), 448-445. <https://doi.org/10.35816/jiskh.v10i2.314>
- Sitanggang, E., Kasim, F., & Sari, N. M. (2020). Analisis Faktor Yang Memengaruhi Kejadian Stunting Di Wilayah Kerja Puskesmas Perdagangan Tahun 2020. *Jurnal Kajian Kesehatan Masyarakat*, 2(1), 141-149. <https://ejournal.delihusada.ac.id/ejournal/index.php/JK2M/article/download/459/309>
- Wiyandani, H. T. (2019). Pelayanan Antenatal Care di Puskesmas. *Higeia Journal of Public Health*, 3(4), 579-587.

**Astri Mufti Aini\***, Maria Komariah, Lilis Mamuroh, Laili Rahayuwati, Yanti Hermayanti, Ristina Mirwanti, Theresia Eriyani

Fakultas Keperawatan Universitas Padjadjaran  
Corresponding author: \*E-mail: [astri20006@mail.unpad.ac.id](mailto:astri20006@mail.unpad.ac.id)

DOI: <https://doi.org/10.33024/minh.v7i2.226>